

# Adolescent RSA Responses During an Anger Discussion Task: Relations to Emotion Regulation and Adjustment

Lixian Cui  
University of Toronto

Amanda Sheffield Morris, Amanda W. Harrist,  
Robert E. Larzelere, and Michael M. Criss  
Oklahoma State University

Benjamin J. Houtberg  
Fuller Theological Seminary

The current study examined associations between adolescent respiratory sinus arrhythmia (RSA) during an angry event discussion task and adolescents' emotion regulation and adjustment. Data were collected from 206 adolescents (10–18 years of age,  $M$  age = 13.37). Electrocardiogram (ECG) and respiration data were collected from adolescents, and RSA values and respiration rates were computed. Adolescents reported on their own emotion regulation, prosocial behavior, and aggressive behavior. Multilevel latent growth modeling was employed to capture RSA responses across time (i.e., linear and quadratic changes; time course approach), and adolescent emotion regulation and adjustment variables were included in the model to test their links to RSA responses. Results indicated that high RSA baseline was associated with more adolescent prosocial behavior. A pattern of initial RSA decreases (RSA suppression) in response to angry event recall and subsequent RSA increases (RSA rebound) were related to better anger and sadness regulation and more prosocial behavior. However, RSA was not significantly linked to adolescent aggressive behavior. We also compared the time course approach with the conventional linear approach and found that the time course approach provided more meaningful and rich information. The implications of adaptive RSA change patterns are discussed.

*Keywords:* respiratory sinus arrhythmia, RSA time course, emotion regulation, prosocial behavior, aggressive behavior

Difficulty managing emotions behaviorally and physiologically plays a critical role in the development of various adjustment problems (Steinberg & Avenevoli, 2000; Thayer, Hansen, Saus-Rose, & Johnsen, 2009), whereas the ability to regulate emotions has been associated with positive adaptation despite various adverse circumstances (Buckner, Mezzacappa, & Beardslee, 2003;

Cicchetti, 2010). Examining factors related to social and emotional development during adolescence is critical because of increases in intense emotions and dynamic fluctuations of emotions that occur during this developmental period (Granic, Hollenstein, Dishion, & Patterson, 2003; Steinberg, 2001; Steinberg & Morris, 2001). In the past few decades, with the development of physiological measurement tools and quantification methods, various studies have investigated the neurophysiological mechanism of cardiac vagal control, focused on respiratory sinus arrhythmia (RSA) as an indicator of parasympathetic regulation linked to both emotion and behavioral regulation, as well as child and adolescent adjustment (e.g., Beauchaine, 2001; Berntson, Cacioppo, & Quigley, 1991; Hastings et al., 2008; Kreibig, 2010; Porges, 2007; Porges et al., 2007). However, the findings from these studies are often inconsistent and sometimes contradictory (De Vries-Bouw et al., 2011; Egizio et al., 2008; Eisenberg et al., 2012; El-Sheikh et al., 2009; Hastings et al., 2008; Obradović, Bush, & Boyce, 2011). In addition, a number of methodological and research design limitations in the literature have been identified (Dennis, Buss, & Hastings, 2012; Obradović et al., 2011; Obradović, Bush, Stamperdahl, Adler, & Boyce, 2010). For example, recent studies have identified the importance of considering the context (e.g., types of tasks, emotional arousal level) of RSA data collection and focused on the temporal changes in RSA rather than examining simple change or static measures from baseline to task (Brooker & Buss, 2010; Miller et al., 2013; Obradović et al., 2011; Vögele, Sorg, Studt-

---

This article was published Online First February 2, 2015.

Lixian Cui, Department of Psychology, University of Toronto; Amanda Sheffield Morris, Amanda W. Harrist, Robert E. Larzelere, and Michael M. Criss, Department of Human Development and Family Science, Oklahoma State University; Benjamin J. Houtberg, Department of Marriage and Family, Fuller Theological Seminary.

This research was supported by a United States Department of Agriculture, Oklahoma Agriculture Experiment Station Project grant (AB-1-13921), an Oklahoma Center for the Advancement of Science and Technology (OCAS) grant (AA-5-40772 and AA-5-45433; Project No. HR11-130), and a Eunice Kennedy Shriver National Institute of Child Health and Human Development R15 grant (AA-5-43382, 1R15HD072463-01).

This investigation is based on a doctoral dissertation submitted to Oklahoma State University by the first author, under the supervision of the second and third authors.

Correspondence concerning this article should be addressed to Lixian Cui, Department of Psychology, University of Toronto, 3359 Mississauga Road North, Mississauga ON L5L 1C6, Canada. E-mail: lx.cui@utoronto.ca

mann, & Weber, 2010). Nevertheless, there have been few published investigations adopting such time course approaches (i.e., temporal changes across time). In response, in the current study we aimed to examine the associations between time course changes in RSA and adolescent emotion regulation and adjustment in a low-income, ethnically diverse sample during an angry event discussion task in a laboratory setting. In addition, we compared the results using the time course approach and those using conventional approach for analyzing RSA data.

### RSA, Baseline RSA, and Adjustment

RSA refers to “a rhythmical fluctuation in heart periods (i.e., interbeat interval) at the respiratory frequency that is characterized by a shortening and lengthening of heart periods in a phase relationship with inspiration and expiration, respectively” (Bernston, Cacioppo, & Quigley, 1993, p. 183). It reflects changes in heart rate, at a high respiratory frequency band (about 0.12–0.40 Hz for adolescents in resting state), which is primarily the function of the parasympathetic nervous system (PNS; the Polyvagal theory, Porges, 1995b, 2011). Tonic parasympathetic control of heart is usually quantified as baseline RSA (Diamond, Fagundes, & Butterworth, 2012; Porges, 1995a) and is assessed when an individual is at rest with paced breathing, or when an individual is shown a calming film (e.g., a film showing dolphins swimming; Fabes & Eisenberg, 1997). It is posited that baseline RSA reflects an individuals’ ability to focus attention, engage in social communication (i.e., social engagement), and maintain homeostasis under normal circumstances (El-Sheikh et al., 2009; Porges, 2007). Baseline RSA also may denote physiological regulation of inner resources, potential for emotion regulation capacity, or a trait-like level of arousability (Beauchaine, 2001; Hastings et al., 2008; Porges, 2007). Empirically, higher baseline RSA has been linked to higher effortful control scores and fewer aggressive behaviors among adolescents (Chapman, Woltering, Lamm, & Lewis, 2010). Likewise, Vögele et al. (2010) found that lower resting high frequency heart rate variability (similar to resting/baseline RSA) was observed among individuals with anger rumination tendencies compared to those with adaptive coping styles. Moreover, lower baseline RSA also has been observed among parasuicidal adolescent girls compared to control group girls (Crowell et al., 2005), and among aggressive adolescents with conduct and attention deficit hyperactivity disorder problems (e.g., Beauchaine, Hong, & Marsh, 2008). A longitudinal study also found higher initial baseline RSA was associated with declines in externalizing symptoms, whereas lower initial baseline RSA was associated with increases in externalizing symptoms among boys (El-Sheikh & Hinnant, 2011).

### RSA Reactivity and Adjustment

RSA reactivity in response to challenge reflects changes in parasympathetic control of the heart from a resting state to a challenging condition and can be quantified as RSA augmentation (i.e., RSA increases from baseline or resting state to challenging condition) or RSA suppression (i.e., RSA decreases; El-Sheikh et al., 2009). RSA suppression indicates parasympathetic or vagal withdrawal, which generates increases in heart rate. RSA augmentation indicates vagal augmentation, which typically slows heart

rate and inhibits sympathetic nervous system input (e.g., El-Sheikh et al., 2009; Obradović & Boyce, 2012). RSA augmentation facilitates the maintenance of an internal equilibrium and social engagement, and RSA suppression signifies the readiness to respond to internal or external threats or challenges (Brooker & Buss, 2010; Porges, 2007). Studies have found that participants with greater RSA suppression during experimental challenges such as star tracing and watching film clips, showed fewer behavioral problems, higher peer status, higher sociability, higher levels of social skills, and better emotion regulation skills during childhood and adolescence (e.g., Beauchaine, Gatzke-Kopp, & Mead, 2007; El-Sheikh, Hinnant, & Erath, 2011; Gentzler, Santucci, Kovacs, & Fox, 2009; see Graziano & Derefinko, 2013, for a review).

However, other studies have found either no relation between RSA reactivity and emotion and behavior problems during childhood and adolescence (e.g., Eisenberg et al., 2012; Hinnant & El-Sheikh, 2009; Whitson & El-Sheikh, 2003) or contradictory results. For example, greater parasympathetic withdrawal in response to public speaking has been associated with higher rates of antisocial behavior reoffending among adolescent males (e.g., De Vries-Bouw et al., 2011). In contrast, less RSA suppression in response to cognitive challenges was related to better social functioning among older adults (Egizio et al., 2008). In addition, higher RSA augmentation in response to a social challenge has been linked to fewer internalizing and externalizing problems and better behavior self-regulation among young children (Hastings et al., 2008). Hinnant and El-Sheikh (2009) did not find significant associations between RSA suppression and internalizing and externalizing problems, but greater RSA suppression in conjunction with low baseline RSA predicting higher internalizing problems from middle childhood to early adolescence. Taken together, generally in RSA reactivity empirical studies, both RSA suppression and augmentation have been observed in response to laboratory challenges. Findings of RSA reactivity in relation to adjustment are mixed (see also Hastings & Miller, 2014, for a review). In addition, we concur with other researchers (e.g., Hastings et al., 2008; Obradović et al., 2011; Overbeek, van Boxtel, & Westerink, 2012; Porges, 2007, 2011) that the association between parasympathetic reactivity and adjustment depends on the type and intensity of specific laboratory challenging task used to quantify RSA reactivity.

### Characteristics of Laboratory Stressors and RSA Reactivity

In RSA empirical studies, the context of measurement and methods used to induce emotion may determine which type of RSA reactivity can be observed (Obradović et al., 2011; Overbeek et al., 2012). In particular, under conditions of high stress such as a public speech, it is more likely to observe RSA suppression, which reflects physiological arousal and stress reactivity, and may indicate an individual’s higher potential and readiness to cope with the stress. Under conditions of mild stress such as a social interview or social activities such as parent–adolescent interactions, it is more likely to observe RSA augmentation or maintenance, which may reflect an individuals’ active regulation to facilitate social engagement/communication (see Porges, 2003, 2007, 2011). Further, the context of measurement and methods used to induce emotion also may determine which of the two types of reactivity is

adaptive or maladaptive (Overbeek et al., 2012). For example, the association between parasympathetic regulation and emotional arousal and coping has been found only under conditions of moderate and high stressors (Fabes & Eisenberg, 1997). Its link to major depressive disorder was significant only in response to a physical stress (squeezing a handgrip dynamometer) but not to cognitive stress (working memory N-back; Nugent, Bain, Thayer, Sollers, & Drevets, 2011). The characteristics of laboratory stressors are critical in psychophysiological studies (Dickerson & Kemeny, 2004) in that various stressors might evoke levels of social and attentional vigilance or alter affective states differentially (Egizio et al., 2008; El-Sheikh, 2005a, 2005b). RSA suppression and augmentation may represent different processes in response to various lab stressors. For example, Kop and colleagues (2011) found greater parasympathetic withdrawal during a high cognitive load Stroop color word test, but much less suppression during mood induction. However, the characteristics of lab stressors have not been discussed in relation to RSA reactivity. Moreover, a large body of extant literature has focused on physical and cognitive stressors, and studies examining emotional stressors typically use emotion (often fear) eliciting pictures and film clips. None of these stressors reflect adolescents' real-life emotion-charged social experience. The current study used a 3-min parent-adolescent emotion (i.e., angry) event discussion task with two phases (i.e., Phase 1: stressful angry event recalling, and Phase 2: regulating anger) and investigated RSA time course changes across the two phases.

### The Study of Emotion Regulation and RSA in Real Time

It is widely accepted that emotion and emotion regulation are rapid processes dependent on contextual demands, with individuals displaying fluctuations and flexibilities in emotion responses to unique situational contexts (e.g., Camras & Witherington, 2005; Thompson, 1994). Cole, Martin, and Dennis (2004) suggested more broadly that emotion regulation is "systematic changes associated with activated emotions" (p. 320). The stress-sensitive physiological regulation systems such as the PNS are either acutely or chronically changing over time within individual in accordance with changing internal and external contexts, which is called *allostasis* (Adam, 2012). However, previous RSA studies are variable centered and typically tap only interindividual differences, and RSA changes are generally averaged across time. RSA reactivity is often computed by simple change scores (subtracting the averaged scores of RSA levels across a challenging task from baseline RSA), or residualized change scores (regressing the averaged RSA values on baseline RSA; we labeled such methods the conventional approach). These scores do not reflect the temporal emotion and emotion regulation processes, which may be the reason for the inconsistent findings in empirical RSA studies. Butler, Wilhelm, and Gross (2006) posited that within-individual changes in RSA are particularly related to regulatory effort. By the same token, we argue that modeling the intraindividual temporal RSA changes over the processes of a specific laboratory task is a better approach (i.e., the time course approach) allowing researchers to examine the adaptability of RSA reactivity in specific laboratory contexts, as the RSA studies involving social and emotional processes increase (e.g., Brooker & Buss, 2010; Burt &

Obradović, 2013; Butler et al., 2006; Miller et al., 2013; Vögele et al., 2010).

To date, some studies have examined PNS regulation processes by specific experimental design such as having a baseline task, a stressful public speech preparation and speech task, and a recovery period (Gunnar, Frenn, Wewerka, & van Ryzin, 2009); a baseline, anger provocation, anger contemplation, and recovery (Vögele et al., 2010); or separating specific components of a task such as breaking up watching a 3-min film clip into 30-s epochs (Crowell et al., 2005). Vögele et al. (2010) found that the pattern of PNS activation in response to anger provocation was observed among anger ruminators, whereas the pattern of PNS activation during anger contemplation and recovery phases occurred among cognitive reappraisers. These findings suggest that PNS activation may be maladaptive if emotional arousal is expected and may be adaptive if the regulation of negative emotions is required.

Researchers also have investigated the temporal RSA changes within individuals and have linked RSA change patterns to adjustment. For example, longitudinally, researchers have modeled changes in RSA reactivity across 3 years among adolescents and found that as RSA change scores from baseline to emotional challenge condition increased across the 3 years, adolescents experienced fewer difficulties with emotion regulation (Vasilev, Crowell, Beauchaine, Mead, & Gatzke-Kopp, 2009). Brooker and Buss (2010) modeled fluctuations (i.e., linear and quadratic changes) in RSA within individual toddlers during a 2½-min "stranger task" using growth curve models, and found that high fear toddlers showed greater linear increases and more negative quadratic change (RSA augmentation followed by RSA suppression) than nonhigh fear children. It was not surprising that the two groups did not differ on simple changes in RSA during the task from baseline. Likewise, Miller and colleagues (2013) used latent growth modeling and captured nonlinear dynamic RSA changes during anger induction, and found that initial RSA suppression and subsequent RSA recovery to baseline was associated with better regulation of aggression among 4- to 6-year-old children, and this pattern of RSA dynamic change also was associated with positive parenting styles. However, less is known about RSA time course changes in response to social and emotional challenges among adolescents, especially adolescents residing in disadvantaged neighborhoods.

### The Current Study

The current study considered the context of measurement and utilized a time course approach to examine RSA reactivity (i.e., modeled both adolescent RSA linear and quadratic changes during the angry event discussion with their parent). In particular, adolescents were asked to talk about a time during the last month when they got angry/mad, while they were away from home and their parents were not around. Adolescents were asked to describe where they were, who they were with, what they were doing, why they got angry, and what they did about it, while parents were asked to listen first and comment or ask questions later. Phase 1 of the task was posited to be quite stressful and may induce quite intensive emotion experience and Phase 2 of the task required regulation during and after the adolescent was recalling the angry event and/or reflected maintaining social engagement in the conversation with parents. An angry task was chosen because anger

and anger regulation may be very important for adolescents, particularly in our sample of youth, living in disadvantaged neighborhoods where adolescents are frequently exposed to violence, gang activity, and peer victimization (Cui, Morris, Criss, Houlberg, & Silk, 2014; Houlberg, Henry, & Morris, 2012; Ingoldsby & Shaw, 2002). We aimed to answer the following questions: (1) What are the overall RSA time course responses during an anger discussion task? (2) How do baseline RSA and the RSA time course changes relate to emotion regulation and adjustment outcomes (i.e., anger and sadness regulation, prosocial, and aggressive behavior)? (3) Does the time course approach provide richer information (e.g., more associations with adjustment outcomes) compared to the conventional approach?

During the angry task, by design, adolescents were expected (1) to be emotionally aroused with anger (due to retrieving the angry memories during Phase 1 of the task, predominantly emotional arousal); and (2) to regulate the corresponding emotions thereafter (due to having the parent's help in dealing with the emotional experience while discussing the event during Phase 2 of the task, predominantly emotion regulation) and/or to maintain engaging in the discussion with parents. Therefore, the minute-to-minute temporal changes in RSA were expected to correspond to processes of adolescent emotional arousal (RSA suppression) and regulation (RSA rebound, increases toward baseline). We hypothesized that overall, we would observe RSA suppression (a negative linear coefficient) initially followed by RSA rebound (a positive quadratic coefficient) from the baseline and across the task (e.g., Brooker & Buss, 2010; Vögele et al., 2010). Based on previous empirical findings, we also hypothesized that higher baseline RSA and a pattern of greater RSA suppression followed by greater RSA rebound would be related to better adolescent anger and sadness regulation, more prosocial behavior, and less aggressive behavior.

Based on the literature, we statistically controlled for several variables. For example, Overbeek et al. (2012) found that RSA reactivity to picture series and film fragments for inducing specific basic emotions differed by sex. Research also has suggested that baseline RSA develops from infancy to early adolescence (El-Sheikh, 2005a, 2005b; Rigterink, Katz, & Hessler, 2010) and contextual factors affect the development of RSA baseline and reactivity (Katz & Rigterink, 2012). Therefore, controlling for adolescents' sex and age in analyses is important. In addition, it is essential to control for respiration rate (Berntson et al., 1997; Berntson, Quigley, & Lozano, 2007; Kok & Fredrickson, 2010) because of the potential influence of respiration, which may be important when youth are talking during discussion tasks with parents. Talking and dramatic physical movements, which alter respiration, may confound RSA measurement (Butler et al., 2006; Ritz, 2009; Tininenko, Measelle, Ablow, & High, 2012).

Most RSA studies have not examined the effects of socioeconomic status (SES) or income and ethnicity. To our knowledge, no study has found that the associations between RSA baseline and reactivity and adjustment differ by SES or ethnicity. Thus, we expected to find similar patterns in these associations in our low-income sample as had been found in other studies with high SES samples. Given that none of the ethnic groups had big enough sample sizes to warrant ethnic group comparisons, such analyses were not done. We also were interested in comparing the time course approach with the conventional approach. Therefore, we averaged RSA values across the task and ran similar models

focusing on linear RSA change slope from baseline to the task overall. We expected that the time course approach would yield richer information compared to the conventional approach because the conventional approach can only tap into simple change.

## Method

### Participants

The sample consisted of 206 families with adolescents who participated in the Family and Youth Development Project (FYDP; Cui et al., 2014). The purpose of the FYDP was to examine predictors and outcomes of adolescent emotion regulation and parent emotional socialization. Data were collected from urban and rural areas of a southern Midwest region of the United States from adolescents ( $M$  age = 13.37 years,  $SD$  = 2.32, range = 10–18 years; 51% female; 32% African American, 29.6% European American, 19.4% Latino American, 19% other) and while interacting with their primary caregivers (83.3% biological mothers, 10.7% biological fathers, 2% grandparents, 4% other primary caregivers). The sample was predominantly comprised of low-income families ( $Mdn$  annual income = \$40,000, 47.5% of families were receiving welfare or public assistance) with an average of 4.35 people living in each home, and 38.7% headed by single parents.

### Procedure

Institutional Review Board (IRB) approval was obtained prior to recruitment and data collection. Participants were recruited from disadvantaged communities with high percentages of ethnic minority and low-income families through fliers distributed at local Boys and Girls Clubs and public facilities, such as the Young Men's Christian Association. A snowball sampling method was also used (had participants give study fliers to friends) although less than 5% of participants were recruited using this method. Interested parent and adolescent dyads were asked to come to a university laboratory to participate in the study. Following the IRB approved procedures, the purpose and procedure of the study were explained to adolescents and their primary caregivers before they signed assent and consent forms, respectively. Next, they were taken to separate rooms where a packet of questionnaires was provided. Research assistants read questions while participants filled in the questionnaires on their own. Parents were taken to the adolescents' room after both finished their questionnaires and completed some interaction tasks.

After completing questionnaires, Biopac MP150 (Biopac Systems, Santa Barbara, CA) cardiovascular (ECG100C) electrodes were attached to the child's shoulder and sides (three-lead) and a respiratory (RSP100C) belt was wrapped around the child's chest while the parent was present. They were then shown a 90-s documentary film clip about dolphins while electrocardiogram (ECG) and respiration signals were recorded to get the baseline RSA (the last 60 s were used). Next, parents and adolescents completed several observational tasks. The interaction task used in this investigation was the angry event discussion task. Instructions for the adolescent were explained by the research assistant and stated,

From time to time, we all feel strong emotions. I'd like you to talk about a time during the last month when you got *angry* when your parents weren't around. Talk with your [mother] about what you did. Go into as much detail as you like, starting from the beginning and going to the end, describing where you were, why you got angry, and what you did about it.

Instructions for parent: "[parent's name], please first listen to [target youth's name] and then comment or ask questions about anything you may be interested in." ECG and respiration signals were recorded during the task, and RSA was calculated minute-to-minute. The laboratory assessment lasted 2 hr on average. The parent and adolescent each received \$60 compensation for their time spent in the lab and were debriefed after the research session was completed.

## Measures

**Adolescent emotion regulation.** Adolescents reported on their abilities to regulate their emotions using the Children's Emotion Management Scale: Sadness and Anger scales (CSMS, Zeman, Shipman, & Penza-Clyve, 2001; CAMS, Zeman, Shipman, & Suveg, 2002). The sadness and anger coping scales were used in this study as indicators of adolescent emotion regulation. The sadness subscale includes five items such as "When I am feeling sad, I control my crying and carrying on" and "I stay calm and don't let sad things get to me." One item ("When I am sad, I do something totally different until I calm down") was deleted to improve reliability (Cronbach's  $\alpha$  was .61 in this study). The anger subscale included four items such as "When I am feeling mad, I control my temper" and "I can stop myself from losing my temper" (Cronbach's  $\alpha$  was .74 in this study). The scale ranged from 0 (*not true*) to 2 (*very true*). Previous studies have demonstrated adequate internal validity of the scales with a Cronbach's alpha ranging from .62 to .77 (Sullivan, Helms, Kliever, & Goodman, 2010; Zeman et al., 2001, 2002). Higher scores indicate better emotion regulation skills.

**Adolescent prosocial behavior.** Adolescents reported on their prosocial behavior during the past year based on a 3-point scale ranging from 0 (*not true*), 1 (*sometimes*), to 2 (*true*). This measure (from the Strengths and Difficulties Questionnaires; SDQ; Goodman & Scott, 1999) included five items such as "I try to be nice to other people, I care about their feelings" and "I usually share with others." Cronbach's alpha was .81 in the current study. Higher scores indicate more prosocial behavior.

**Adolescent aggressive behavior.** Adolescent aggressive behavior was obtained using 14 items from the Problem Behavior Frequency Scale (Farrell, Kung, White, & Valois, 2000), which assessed the frequency of physical, relational, and verbal aggression, such as "get in a fight in which someone was hit," "spread a rumor," and "insult someone's family." Adolescents were asked to indicate how frequently they engaged in each behavior during the past year using the following scale ranging from 1 (*never*), 2 (*1–2 times*), 3 (*3–4 times*), 4 (*5–6 times*), 5 (*7 or more times*). Cronbach's alpha was .88 in the current study. Higher scores indicate more aggressive behavior.

**RSA and respiration rate.** Adolescent ECG and respiration pneumogram were recorded using Biopac MP150 system. The sampling frequency was 2 kHz for both ECG and respiration signals. ECG and respiration data were imported to MindWare

HRV 3.0.9 (MindWare Technologies, Ltd., 2010) to calculate RSA values and respiration rate (e.g., Rash & Prkachin, 2013). We used power spectral analysis method and employed Hamming spectral window in calculating RSA for our short-term recordings of ECG (Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology, [Task Force], 1996). RSA values in  $\text{ms}^2$  and respiration rate were calculated during the neutral film baseline task and the 3-min angry task. ECG data were analyzed minute by minute for high frequency component of HRV (Task Force, 1996) and ECG artifacts were edited (including visual checks and manual corrections) in the MindWare HRV 3.0.9 program to obtain reliable measures of RSA. In data preparation and analyses, we controlled for respiration in two ways: (1) computing respiration peak frequency in MindWare HRV 3.0.9, we discarded unreliable RSA values when respiration frequency was out of the high frequency 0.12 to 0.40 Hz range (Berntson, Norman, Hawkey, & Cacioppo, 2008); (2) including respiration rates in Level 1 of multilevel models as covariates. Twenty-two adolescents' physiological data during the discussion task were not usable due to technical issues or research assistant errors and unreliable RSA values were treated as randomly missing across the 3 min (about 14%). Missing data were handled in HLM 7 (Raudenbush, Bryk, & Congdon, 2011) using full information maximum likelihood (FIML) estimation during multilevel modeling.

## Results

### Descriptive Statistics

Means and standard deviations of all study variables are presented in Table 1. As indicated in Table 1, the zero-order bivariate correlations showed that during the baseline and the second minute of the anger task, higher respiration rate was significantly associated with lower RSA levels, suggesting that respiratory activities may influence RSA and we should control for respiration rate in statistical analyses. Neither respiration rate nor RSA was significantly related to adolescent sex or age (Tables 1). RSA levels were correlated with each other across baseline and the 3-min anger task, but were not significantly associated with adolescent emotion regulation, prosocial behavior, or aggressive behavior. Adolescent sadness and anger regulation abilities were positively associated with prosocial behavior, and negatively related to aggressive behavior. Adolescent prosocial behavior was negatively associated with aggressive behavior. Adolescent age was positively related to aggressive behavior (see Table 1). Finally, adolescent girls had higher levels of prosocial behavior,  $t(204) = 4.30, p < .001$ , and lower levels of aggressive behavior than boys,  $t(204) = -1.99, p = .048$ .

### RSA Reactivity in Response to Anger Event Discussion

**The time course approach.** To answer the first research question about the overall RSA time course changes across time, we first implemented the time course approach using multilevel modeling in HLM 7 by incorporating both linear and quadratic effects of time (across all four time points, 1-min baseline and 3-min task) while controlling for respiration rate, so that the

Table 1  
Means, Standard Deviations, and Correlations of Major Variables

	<i>N</i>	<i>M</i>	<i>SD</i>	1	2	3	4	5	6	7	8	9	10	11	12	13
1. Sex <sup>a</sup>	206	0.49	0.50	—												
2. Age	206	13.37	2.32	-.01	—											
3. RSA baseline	188	7.09	1.32	-.03	-.12	—										
4. RR baseline	188	18.70	4.99	-.04	-.02	-.15*	—									
5. RSA M1	155	7.01	1.17	.03	-.10	.65***	.04	—								
6. RR M1	155	21.64	6.28	-.03	-.01	-.04	.08	-.05	—							
7. RSA M2	155	6.90	1.23	.10	-.07	.73***	-.04	.81***	-.00	—						
8. RR M2	155	21.63	5.37	.01	-.03	-.01	.03	-.03	.12	-.16*	—					
9. RSA M3	154	6.95	1.10	.05	-.11	.64***	.11	.77***	-.08	.78***	-.13	—				
10. RR M3	154	21.50	5.59	-.07	-.12	-.02	.17*	-.00	.19*	.04	.27***	.04	—			
11. SR	205	1.26	0.47	-.01	.02	.03	-.06	.06	.05	.03	.04	.10	-.12	—		
12. AR	206	1.19	0.52	-.02	-.03	-.00	.00	-.04	.04	-.05	.02	.07	.05	.40***	—	
13. Prosocial	206	1.57	0.43	-.29***	-.06	.14 <sup>†</sup>	-.02	-.03	.11	.01	.02	.10	.04	.28***	.45***	—
14. AB	206	1.43	0.52	.14*	.18**	-.08	.09	-.09	-.16*	-.09	.03	-.03	-.12 <sup>†</sup>	-.43***	-.28***	—

Note. RSA = respiratory sinus arrhythmia; RR = respiration rate; M1 = first minute of anger task; M2 = second minute of anger task; M3 = third minute of anger task; SR = sadness regulation; AR = anger regulation; AB = aggressive behavior.

<sup>a</sup> Adolescent sex was coded as 0 (female) and 1 (male).

<sup>†</sup>  $p < .10$ . \*  $p < .05$ . \*\*  $p < .01$ . \*\*\*  $p < .001$ .

coefficients of linear and quadratic terms represent RSA reactivity. At second level, adolescent sex and age were controlled. The results of this model without further predictors showed that there was a significant initial decrease in RSA,  $b_2 = -0.44, p < .001$ , followed by a significant increase in RSA,  $b_3 = 0.10, p = .002$ , indicating a pattern of initial RSA suppression and subsequent RSA rebound. No age or sex differences in RSA baseline, respiration effects, RSA suppression, and rebound were found.

**The conventional approach.** Next, we implemented the conventional approach by computing a single slope (linear change) from baseline to the discussion task overall. RSA reactivity was then the mean RSA change from the baseline to the 3-min task overall. In the multilevel model, time was dummy coded (0 for the RSA baseline; 1 for each minute of the anger task), so that the intercept represents RSA baseline (time = 0), and the coefficient for time represents the mean change from baseline to the anger task. Likewise, adolescent concurrent respiration rates were controlled for at Level 1 and adolescent age and sex were controlled for at Level 2. The results of this model showed that there was a significant decrease in adolescent RSA from baseline to the anger task,  $b_2 = -0.42, p = .001$ , indicating significant RSA suppression from baseline to the anger task. No age or sex differences were found.

**Baseline RSA, RSA Reactivity, and Emotion Regulation and Adjustment<sup>1</sup>**

**The time course approach.** To answer the second research question regarding the associations between baseline RSA, RSA reactivity, and adolescent emotion regulation and adjustment, based on the previous model, adolescent anger and sadness regulation, and prosocial and aggressive behavior were entered separately to predict RSA baseline ( $b_0$ ), and RSA linear change ( $b_2$ ) and quadratic change ( $b_3$ ) in the following model (four separate models were tested):

$$\text{Level 1: } RSA = b_0 + b_1 RR + b_2 \text{Time} + b_3 \text{Time}^2 + \varepsilon.$$

$$\text{Level 2: } b_0 = \gamma_{00} + \gamma_{01} \text{Predictor} + \gamma_{02} \text{Sex} + \gamma_{03} \text{Age} + v_0.$$

$$b_1 = \gamma_{10} + \gamma_{11} \text{Predictor} + \gamma_{12} \text{Sex} + \gamma_{13} \text{Age} + (v_1).$$

$$b_2 = \gamma_{20} + \gamma_{21} \text{Predictor} + \gamma_{22} \text{Sex} + \gamma_{23} \text{Age} + v_2.$$

$$b_3 = \gamma_{30} + \gamma_{31} \text{Predictor} + \gamma_{32} \text{Sex} + \gamma_{33} \text{Age} + v_3.$$

Respiration rates (RR) were grand mean centered at Level 1. Time was coded as 0 (RSA baseline) and 1, 2, 3 (first-, second-, and third-minute RSA during the anger task, respectively). Predictors (i.e., sadness and anger regulation, prosocial and aggressive behavior) were tested separately. Predictor and adolescent age and sex were grand mean centered at Level 2. Error term in parentheses ( $v_1$ ) was trimmed in final models because it was not significant.

As shown in Table 2, the results showed that adolescents with high sadness regulation had greater initial RSA suppression,  $\gamma_{21} = -0.48, p = .04$ , and greater subsequent RSA rebound,  $\gamma_{31} = 0.15, p = .03$  (see Figure 1). Likewise, adolescents with high anger regulation showed greater initial RSA suppression,  $\gamma_{21} = -0.74, p < .001$ , and greater subsequent RSA rebound,  $\gamma_{31} = 0.20, p < .001$  (see Figure 2). Adolescents with high anger regulation also demonstrated marginally high baseline RSA,  $\gamma_{01} = 0.33, p = .08$ . Adolescents with high prosocial behavior showed high RSA baseline,  $\gamma_{01} = 0.68, p = .01$ , greater initial RSA suppression,  $\gamma_{21} = -1.02, p < .001$ , and greater subsequent RSA rebound,  $\gamma_{31} = 0.30, p < .001$  (see Figure 3). We

<sup>1</sup> Parent-reported adolescent emotion regulation and adjustment data also were collected and tested in relation to RSA variables but no significant effects were identified. This suggests that adolescents' physiological regulation was related to their own perceptions of emotion regulation and adjustment, but not to their parents' perceptions. It should be noted that there is growing recognition that discrepancies in findings across reporters is not a flaw or limitation. Rather the discrepancies provide essential insight into the nuances and contextualization of our understanding of an individual's regulation functioning and development within complex social relationships involving multiple perspectives.

Table 2  
Four Multilevel Models of Whether Dynamic RSA Changes Were Associated With Adolescent Emotion Regulation and Adjustment (Level 2 Results)

Fixed effect	Model 1 sadness regulation			Model 2 anger regulation			Model 3 prosocial behavior			Model 4 aggressive behavior		
	Coefficient	SE	t	Coefficient	SE	t	Coefficient	SE	t	Coefficient	SE	t
<i>b</i> <sub>0</sub> (RSA baseline)												
γ <sub>00</sub> (intercept)	7.23***	0.11	67.99	7.24***	0.10	69.35	7.25***	0.10	69.52	7.27***	0.11	66.59
γ <sub>01</sub> (predictor)	0.30	0.19	1.57	0.33 <sup>†</sup>	0.18	1.79	0.68**	0.24	1.99	-0.33	0.22	-1.47
γ <sub>02</sub> (sex)	-0.00	0.21	-0.01	-0.02	0.21	-0.11	0.17	0.21	0.81	0.00	0.21	0.02
γ <sub>03</sub> (age)	-0.08	0.05	-1.65	-0.08 <sup>†</sup>	0.04	-1.74	-0.07	0.04	-1.69	-0.05	0.05	-1.04
<i>b</i> <sub>1</sub> (respiration rate)												
γ <sub>10</sub> (intercept)	-0.01*	0.01	-2.13	-0.01**	0.01	-2.53	-0.02*	0.01	-2.37	-0.02*	0.01	-2.57
γ <sub>11</sub> (predictor)	-0.02 <sup>†</sup>	0.01	-1.94	-0.03**	0.01	-3.27	-0.03*	0.01	-2.35	0.02	0.01	1.53
γ <sub>12</sub> (sex)	-0.01	0.01	-0.52	-0.00	0.01	-0.40	-0.01	0.01	-1.15	-0.01	0.01	-0.44
γ <sub>13</sub> (age)	0.00	0.00	0.46	0.00	0.00	0.55	0.00	0.00	0.51	-0.00	0.00	-0.08
<i>b</i> <sub>2</sub> (RSA linear change)												
γ <sub>20</sub> (intercept)	-0.38**	0.12	-3.21	-0.40***	0.11	-3.56	-0.43***	0.12	-3.67	-0.44**	0.13	-3.48
γ <sub>21</sub> (predictor)	-0.48*	0.24	-2.04	-0.74**	0.17	-4.32	-1.02*	0.27	-3.82	0.16	0.27	0.60
γ <sub>22</sub> (sex)	0.18	0.24	0.74	0.21	0.23	0.92	-0.10	0.25	-0.40	0.24	0.25	0.97
γ <sub>23</sub> (age)	0.04	0.05	0.80	0.05	0.05	0.91	0.04	0.05	0.82	0.03	0.06	0.44
<i>b</i> <sub>3</sub> (RSA quadratic change)												
γ <sub>30</sub> (intercept)	0.09**	0.03	2.77	0.10**	0.03	3.09	0.10**	0.03	3.26	0.10**	0.03	3.04
γ <sub>31</sub> (predictor)	0.15*	0.07	2.19	0.20**	0.05	4.35	0.30***	0.07	4.03	-0.03	0.08	-0.42
γ <sub>32</sub> (sex)	-0.05	0.06	-0.76	-0.06	0.06	-0.93	0.03	0.07	0.47	-0.07	0.07	-1.02
γ <sub>33</sub> (age)	-0.01	0.01	-0.62	-0.01	0.01	-0.72	-0.01	0.01	-0.58	-0.01	0.02	-0.38
Random effect	Variance component	χ <sup>2</sup> (df)	p	Variance component	χ <sup>2</sup> (df)	p	Variance component	χ <sup>2</sup> (df)	p	Variance component	χ <sup>2</sup> (df)	p
υ <sub>0</sub> (RSA baseline)	1.35	860.69 (159)	.000	1.36	874.41 (159)	.000	1.33	876.90 (159)	.000	1.34	846.59 (159)	.000
υ <sub>2</sub> (RSA linear change)	0.40	234.16 (159)	.000	0.38	231.10 (159)	.000	0.39	233.26 (159)	.000	0.39	231.86 (159)	.000
υ <sub>3</sub> (RSA quadratic change)	0.02	202.27 (159)	.012	0.02	200.69 (159)	.014	0.02	200.24 (159)	.015	0.02	202.18 (159)	.012
Level 1 error, ε <sub><i>t</i></sub>	0.32			0.32			0.32			0.32		

Note. RSA = respiratory sinus arrhythmia.  
<sup>†</sup> *p* < .10. \* *p* < .05. \*\* *p* < .01. \*\*\* *p* < .001.

did not find significant results for aggressive behavior (see Table 2). Taken together, these results suggest that a pattern of initial RSA suppression and subsequent RSA rebound was associated with better sadness and anger regulation, and more

prosocial behavior. Further, higher RSA baseline was related to more prosocial behavior.

**The conventional approach.** Next, we again implemented the conventional approach, similarly based on the previous




Figure 1. Respiratory sinus arrhythmia (RSA) changes across time for adolescents with different levels of sadness regulation. SD = standard deviation; M = minute. See the online article for the color version of this figure.




Figure 2. Respiratory sinus arrhythmia (RSA) changes across time for adolescents with different levels of anger regulation. SD = standard deviation; M = minute. See the online article for the color version of this figure.




Figure 3. Respiratory sinus arrhythmia (RSA) changes across time for adolescents with different levels of prosocial behavior. *SD* = standard deviation; M = minute. See the online article for the color version of this figure.

model, adolescent anger and sadness regulation, and adolescent prosocial and aggressive behavior scores were entered at Level 2 separately to predict baseline RSA ( $b_0$ ) and RSA suppression ( $b_2$ ) in the following model (four separate models were tested):

$$\text{Level 1: } RSA = b_0 + b_1 RR + b_2 \text{Time} + \varepsilon.$$

$$\text{Level 2: } b_0 = \gamma_{00} + \gamma_{01} \text{Predictor} + \gamma_{02} \text{Sex} + \gamma_{03} \text{Age} + v_0.$$

$$b_1 = \gamma_{10} + \gamma_{11} \text{Predictor} + \gamma_{12} \text{Sex} + \gamma_{13} \text{Age} + v_1$$

$$b_2 = \gamma_{20} + \gamma_{21} \text{Predictor} + \gamma_{22} \text{Sex} + \gamma_{23} \text{Age} + v_2.$$

Respiration rates (*RR*) were grand mean centered at Level 1. Time was coded as 0 at RSA baseline and as 1 during each of the other 3 min. Predictors including sadness and anger regulation, prosocial and aggressive behaviors were tested separately, one at a time. Predictor and adolescent age and sex were grand mean centered at Level 2. Error term in parentheses ( $v_1$ ) was trimmed in final models because it was not significant. The results indicated that adolescents with high anger regulation showed greater RSA suppression from baseline to the anger task,  $\gamma_{21} = -0.65, p = .001$ , also marginally high baseline RSA,  $\gamma_{01} = 0.41, p = .05$ . The other three predictors were not significantly linked to either RSA baseline or RSA suppression. Adolescents with high prosocial behavior demonstrated high RSA baseline,  $\gamma_{01} = 0.77, p = .004$ , greater RSA suppression,  $\gamma_{21} = -0.78, p = .001$ .

### Discussion

To shed light on methodological issues in empirical RSA research, the current study adopted a process view of physiological regulation of emotion, and focused on adolescent RSA time course changes during an interpersonal challenging context. We also controlled for respiration parameters, and adolescent sex and age. Our findings indicated that an adaptive RSA change pattern was initial RSA suppression and subsequent RSA rebound, which was related to better anger and sadness regulation and more prosocial behavior among adolescents. These findings suggest that in response to emotional challenge, a pattern of parasympathetic with-

drawal and subsequent parasympathetic activation may be more adaptive, which may facilitate better adolescent emotion regulation skills and positive development. We also compared the time course approach of capturing RSA linear and quadratic changes with the conventional approach of using simple RSA reactivity scores. We found that the time course approach yielded richer information by adding a rebound component of RSA change and more relations to emotion regulation outcomes, indicating that using simple RSA reactivity scores may be one main reason for null or inconsistent findings of previous RSA empirical studies (Burt & Obradović, 2013; Graziano & Derefinko, 2013; Hastings & Miller, 2014).

### RSA Responses and Emotion Regulation and Adjustment

Adopting a time course approach to model RSA temporal changes during an emotion event discussion task, we found a pattern of initial RSA suppression followed by RSA rebound from baseline across the task. Further, greater RSA suppression and rebound were associated with better anger and sadness regulation skills and more prosocial behavior. These results suggest that in the context of interpersonal challenge, such as talking about an actual anger-charged event, a pattern of initial RSA suppression followed by RSA rebound may reflect adaptive emotion regulation abilities, which in turn, may be related to more positive behavior among adolescents. These findings support the notion that RSA suppression in response to challenges may imply a readiness to respond to social stress (Brooker & Buss, 2010; Porges, 2007). Findings are also in line with recent empirical studies that found adaptive function of RSA suppression (e.g., Beauchaine et al., 2007; El-Sheikh et al., 2011; Gentzler et al., 2009). Further, these results are consistent with other investigations using similar time course approaches and support the notion that initial RSA suppression and subsequent RSA increases may be adaptive RSA reactivity in response to social and emotional challenges (e.g., Brooker & Buss, 2010; Miller et al., 2013). In adolescents' daily life, RSA suppression observed when individuals are exposed to stressful or challenging social environment may be advantageous as adolescents are probably emotionally aroused and RSA suppression allows them to react and respond to stressors. In addition, subsequent RSA rebound observed thereafter may be adaptive as individuals are probably actively regulating their intense negative emotions and trying to maintain a calming state to engage in the social activities (Hastings et al., 2008; Porges, 2007; Vögele et al., 2010).

Following the conventional approach of focusing on the overall change from baseline to the anger task, we found significant RSA suppression from baseline to the task overall, and significant associations between RSA suppression and anger regulation and prosocial behavior. Compared to the conventional approach of analyzing RSA data, the time course approach added a rebound component to RSA changes across time, which better matched with actual behavioral and emotional processes (Burt & Obradović, 2013) and provided richer information as it revealed meaningful associations to anger regulation, prosocial behavior, and also sadness regulation. The time course approach is an emerging approach in psychophysiological studies that researchers have started to utilize, and our findings are consistent with previous investigations utilizing similar temporal approaches and suggest

the promise of such methods (e.g., Brooker & Buss, 2010; Gunnar et al., 2009; Miller et al., 2013).

This time course view of RSA suppression and rebound also aligns with a process view of emotion regulation. *Emotion regulation* has been defined in the literature as the internal and external processes by which the individual manages the occurrence, intensity, and expression of emotions to reach goals or situational demands (Eisenberg & Morris, 2003; Thompson, 1994). Cole et al. (2004) argued that researchers must examine emotion regulation in the context of emotional arousal where emotions are activated for there to be evidence of emotion regulation. Likewise, it may be essential for researchers to examine physiological regulation of emotion in process and in the context of emotional arousal for RSA suppression to be observed. Subsequent RSA rebound likely indicates emotion down regulation of the arousal.

Our findings also highlight the importance of considering the context of RSA measurement (Obradović et al., 2011). It is possible that the degree to which RSA reactivity to stress is adaptive depends on the context in which RSA is assessed (Obradović, 2012). We used an anger event recall and discussion task and examined RSA processes across the discussion. In such a social and emotional challenge context, individuals may be relatively emotionally aroused resulting in sympathetic activation and parasympathetic inhibition that would allow RSA suppression to be observed (El-Sheikh et al., 2009; Obradović & Boyce, 2012). This was what we found for the first phase of the task as adolescents were generally recalling and reliving the anger event. During this phase, RSA suppression is adaptive as the youth are emotionally aroused and getting ready to cope or deal with the negative emotion. The subsequent RSA rebound by the end of the anger event conversation in this study may reflect adolescents' regulation efforts while they were responding to parents' questions and engaging in the calming conversation. RSA rebound is adaptive in the second phase of the task. By considering RSA change processes, corresponding to internal and external needs of emotion regulation, our data afford critical information on the adaptability of RSA reactivity.

Note that we did not find significant associations between the pattern of RSA suppression and rebound and adolescent aggressive behavior. Instead, links to prosocial behavior were evident. According to Polyvagal theory (Porges, 2007), RSA suppression reflects an individual's responses to threatening and challenging stimuli, and RSA augmentation is an indicator of active regulation to support calm engagement and form social bonds (Porges, 2011). Thus, the parasympathetic regulation of emotion may facilitate an individual's proficiency for cooperative, caring, and helping behaviors during social interactions (Hastings & Miller, 2014). For our assessment of parasympathetic activity, it may be more likely that RSA is linked to prosocial behavior rather than aggressive behavior. Indeed, studies have found greater RSA suppression in response to challenges associated with more empathic concern and prosocial behavior (e.g., Graziano, Keane, & Calkins, 2007; Hastings & Miller, 2014; Liew et al., 2011). Our lack of findings for aggression is in contrast to findings of Miller et al.'s (2013) study among young children who used an anger-induction film watching approach (i.e., observing other's anger). These authors found initial RSA suppression followed by RSA recovery pattern was related to children's better control of aggression. It is possible that how children react to others' anger (compared to their own anger)

may be more strongly linked to aggression as indicated by the results of Miller et al.'s study. In addition, it should be noted that their temporal RSA changes were assessed across neutral, anger, and positive emotion film watching, and their measure of control of aggression may have tapped anger regulation. Another possible reason of our lack of findings may be that during the second phase of the anger event discussion, adolescents may choose to talk with their parents rather than attempt to self-regulate. As such, the RSA rebound may not have been sufficiently robust to be linked to aggression.

### RSA Baseline and Emotion Regulation and Adjustment

We also found that adolescent baseline RSA, measured when watching a neutral film clip, was linked to positive development (i.e., prosocial behavior) and (at trend level) to better anger regulation. Higher baseline RSA may reflect emotion regulation potentiality, and it has been linked to fewer behavior problems, such as internalizing and externalizing problems (Beauchaine et al., 2008; Crowell et al., 2005; El-Sheikh & Hinnant, 2011). Our findings add to the empirical evidence that higher baseline RSA is associated with positive developmental outcomes. However, the association between RSA baseline and emotion regulation was relatively weak. Several other studies of baseline RSA have not found significant correlates of vagal tone or baseline RSA with child and adolescent adjustment (e.g., Diamond et al., 2012; Egizio et al., 2008; El-Sheikh et al., 2009; Hinnant & El-Sheikh, 2009). Our findings of RSA baseline emerged after controlling for respiration rate, adolescent sex and age.

Despite our significant findings using a time course approach, the effects of RSA baseline were still relatively weak in the current study. It is possible that laboratory baseline measures do not accurately capture the actual baseline RSA of each individual. In particular, the laboratory setting may be novel and thus be moderately stressful to some participants, despite the fact that they typically have spent some time in the laboratory before the baseline measure. Another reason for a lack of findings may be because our baseline measure was relatively short (only 1 min) and it was measured while adolescents were passively watching a neutral film clip. Third, there may be interaction effects between baseline RSA and the environmental contexts in which adolescents live. Some empirical studies support the notion that high baseline RSA may buffer the negative effects of adverse contexts, instead of manifesting main effects (e.g., El-Sheikh & Hinnant, 2011; El-Sheikh et al., 2011). Multiple and longer baselines under different conditions such as pure resting, paced breathing, and in supine position should be explored in future studies. More important, a more comparable talking baseline may work as a better baseline (see Tininenko et al., 2012).

### Strengths and Limitations

One of the strengths of the current study was that we modeled both the linear and quadratic RSA changes across a challenging task, which allowed us to capture the processes of emotional arousal and regulation that correspond to RSA reactivity. A second strength was that we accounted for respiration parameters (e.g., respiration rate) in all analyses. RSA is usually confounded with

respiratory parameters, and previous studies typically have not corrected the respiratory confounds. Controlling for respiration parameters allowed us to obtain a better measure of RSA, particularly in our study in which adolescents were talking from time-to-time and had some physical movement while sitting in the chair. A third strength was that we focused on interpersonal challenging social settings (i.e., parent–adolescent discussion of a real anger event), which represent typical adolescent daily stressors and may demonstrate strong ecological validity. In previous RSA reactivity studies, participants have been passively exposed to lab stressors such as physical (e.g., receiving special stimuli such as lemon juice on the tongue or squeezing handgrip dynamometer), cognitive (e.g., tracing stars or digit span recitation), social (e.g., structured interview), or emotional (e.g., watching film clips) challenges (Butler et al., 2006; Chapman et al., 2010; Ellis, Essex, & Boyce, 2005; El-Sheikh et al., 2011; Kreibig, 2010; Nugent et al., 2011; Obradović et al., 2010). However, in “real life,” most of the everyday stress that makes children and adolescents vulnerable to adjustment problems are related to social issues. In the current study, we used an angry event discussion task in which participants played an active role in choosing the topic and leading the discussion. Another strength was that our sample was a particularly low-income ethnically diverse sample as there have been few published RSA reactivity studies utilizing such samples. However, caution should be taken when generalizing our findings to other populations.

We also had some limitations in the current study. One limitation was that the baseline measure of RSA was not immediately preceding the anger task and the anger task did not have a specific clear recovery period by design. Although it may not have influenced the current findings, it would have been ideal to have a baseline right before the challenging task. A second limitation was that we only focused on measures of parasympathetic nervous system functioning and did not have pure measures of sympathetic nervous system functioning such as preejection period. PNS functioning has been found to be inconsistent across situations (e.g., Bertsch, Hagemann, Naumann, Schächinger, & Schulz, 2012) and is not always in concordance with reactivity in other physiological systems (e.g., Bauer, Quas, & Boyce, 2002; Cacioppo, Uchino, & Berntson, 1994). It would be helpful to investigate various systems simultaneously to better understand physiological response to stress in the future. Recent calls for integrative approaches to studying physiology and emotion across autonomic nervous systems, across multiple biological systems, and even across biological systems and contexts (El-Sheikh et al., 2009; Hastings, Buss, & Dennis, 2012; Kreibig, 2010) point out the need to consider the functioning of different systems. It may be beneficial for future studies that only focused on one physiological system to adopt a process perspective in data analyses and indeed, our findings suggest the advantages of the time course approach. A third limitation was that emotion regulation and adjustment variables were all self-reports. It is possible that adolescents with more adaptive RSA responses report more positive adjustment due to differences in physiological reactivity rather than actual behaviors. Future work should explore such issues using longitudinal data and other outcome measures.

## Conclusions and Implications

The current study findings suggest that a pattern of initial RSA suppression followed by RSA rebound may be adaptive in response to interpersonal stress, especially to promote positive development among adolescents. Methodologically, the current study contributes to the field of RSA research by adopting a time course approach to illustrate the temporal processes of physiological regulation of emotion. The new generation of research on physiology and emotion would benefit from treating physiological regulation of emotion as processes. In addition, much can be gained from the use of multilevel modeling analysis to better understand the complex patterns of physiology of emotion.

## References

- Adam, E. K. (2012). Emotion-cortisol transactions occur over multiple scales in development: Implications for research on emotion and the development of emotional disorders. *Monographs of the Society for Research in Child Development*, 77(2), 17–27. <http://dx.doi.org/10.1111/j.1540-5834.2012.00657.x>
- Bauer, A. M., Quas, J. A., & Boyce, W. T. (2002). Associations between physiological reactivity and children's behavior: Advantages of a multisystem approach. *Journal of Developmental and Behavioral Pediatrics*, 23, 102–113. <http://dx.doi.org/10.1097/00004703-200204000-00007>
- Beauchaine, T. (2001). Vagal tone, development, and Gray's motivational theory: Toward an integrated model of autonomic nervous system functioning in psychopathology. *Development and Psychopathology*, 13, 183–214. <http://dx.doi.org/10.1017/S0954579401002012>
- Beauchaine, T. P., Gatzke-Kopp, L., & Mead, H. K. (2007). Polyvagal theory and developmental psychopathology: Emotion dysregulation and conduct problems from preschool to adolescence. *Biological Psychology*, 74, 174–184. <http://dx.doi.org/10.1016/j.biopsycho.2005.08.008>
- Beauchaine, T. P., Hong, J., & Marsh, P. (2008). Sex differences in autonomic correlates of conduct problems and aggression. *Journal of the American Academy of Child and Adolescent Psychiatry*, 47, 788–796. <http://dx.doi.org/10.1097/CHI.Ob013e318172ef4b>
- Berntson, G. G., Bigger, J. T., Jr., Eckberg, D. L., Grossman, P., Kaufmann, P. G., Malik, M., . . . van der Molen, M. W. (1997). Heart rate variability: Origins, methods, and interpretive caveats. *Psychophysiology*, 34, 623–648. <http://dx.doi.org/10.1111/j.1469-8986.1997.tb02140.x>
- Berntson, G. G., Cacioppo, J. T., & Quigley, K. S. (1991). Autonomic determinism: The modes of autonomic control, the doctrine of autonomic space, and the laws of autonomic constraint. *Psychological Review*, 98, 459–487. <http://dx.doi.org/10.1037/0033-295X.98.4.459>
- Berntson, G. G., Cacioppo, J. T., & Quigley, K. S. (1993). Respiratory sinus arrhythmia: Autonomic origins, physiological mechanisms, and psychophysiological implications. *Psychophysiology*, 30, 183–196. <http://dx.doi.org/10.1111/j.1469-8986.1993.tb01731.x>
- Berntson, G. G., Norman, G. J., Hawley, L. C., & Cacioppo, J. T. (2008). Cardiac autonomic balance versus cardiac regulatory capacity. *Psychophysiology*, 45, 643–652. <http://dx.doi.org/10.1111/j.1469-8986.2008.00652.x>
- Berntson, G. G., Quigley, K. S., & Lozano, D. (2007). Cardiovascular psychophysiology. In J. T. Cacioppo, L. G. Tassinary, & G. G. Berntson (Eds.), *Handbook of psychophysiology* (pp. 182–210). New York, NY: Cambridge University Press. <http://dx.doi.org/10.1017/CBO9780511546396.008>
- Bertsch, K., Hagemann, D., Naumann, E., Schächinger, H., & Schulz, A. (2012). Stability of heart rate variability indices reflecting parasympathetic activity. *Psychophysiology*, 49, 672–682. <http://dx.doi.org/10.1111/j.1469-8986.2011.01341.x>

- Brooker, R. J., & Buss, K. A. (2010). Dynamic measures of RSA predict distress and regulation in toddlers. *Developmental Psychobiology*, *52*, 372–382. <http://dx.doi.org/10.1002/dev.20432>
- Buckner, J. C., Mezzacappa, E., & Beardslee, W. R. (2003). Characteristics of resilient youths living in poverty: The role of self-regulatory processes. *Development and Psychopathology*, *15*, 139–162. <http://dx.doi.org/10.1017/S0954579403000087>
- Burt, K. B., & Obradović, J. (2013). The construct of psychophysiological reactivity: Statistical and psychometric issues. *Developmental Review*, *33*, 29–57. <http://dx.doi.org/10.1016/j.dr.2012.10.002>
- Butler, E. A., Wilhelm, F. H., & Gross, J. J. (2006). Respiratory sinus arrhythmia, emotion, and emotion regulation during social interaction. *Psychophysiology*, *43*, 612–622. <http://dx.doi.org/10.1111/j.1469-8986.2006.00467.x>
- Cacioppo, J. T., Uchino, B. N., & Berntson, G. G. (1994). Individual differences in the autonomic origins of heart rate reactivity: The psychometrics of respiratory sinus arrhythmia and preejection period. *Psychophysiology*, *31*, 412–419. <http://dx.doi.org/10.1111/j.1469-8986.1994.tb02449.x>
- Camras, L. A., & Witherington, D. C. (2005). Dynamic systems approaches to emotional development. *Developmental Review*, *25*, 328–350. <http://dx.doi.org/10.1016/j.dr.2005.10.002>
- Chapman, H. A., Woltering, S., Lamm, C., & Lewis, M. D. (2010). Hearts and minds: Coordination of neurocognitive and cardiovascular regulation in children and adolescents. *Biological Psychology*, *84*, 296–303. <http://dx.doi.org/10.1016/j.biopsycho.2010.03.001>
- Cicchetti, D. (2010). Resilience under conditions of extreme stress: A multilevel perspective. *World Psychiatry*, *9*, 145–154.
- Cole, P. M., Martin, S. E., & Dennis, T. A. (2004). Emotion regulation as a scientific construct: Methodological challenges and directions for child development research. *Child Development*, *75*, 317–333. <http://dx.doi.org/10.1111/j.1467-8624.2004.00673.x>
- Crowell, S. E., Beauchaine, T. P., McCauley, E., Smith, C. J., Stevens, A. L., & Sylvers, P. (2005). Psychological, autonomic, and serotonergic correlates of parasuicide among adolescent girls. *Development and Psychopathology*, *17*, 1105–1127. <http://dx.doi.org/10.1017/S0954579405050522>
- Cui, L., Morris, A. S., Criss, M. M., Houlberg, B. J., & Silk, J. S. (2014). Parental psychological control and adolescent adjustment: The role of adolescent emotion regulation. *Parenting: Science and Practice*, *14*, 47–67. <http://dx.doi.org/10.1080/15295192.2014.880018>
- Dennis, T. A., Buss, K. A., & Hastings, P. D. (2012). Introduction to the monograph. *Monographs of the Society for Research in Child Development*, *77*(2), 1–5. <http://dx.doi.org/10.1111/j.1540-5834.2011.00654.x>
- De Vries-Bouw, M., Popma, A., Vermeiren, R., Doreleijers, T. A., Van De Ven, P. M., & Jansen, L. M. (2011). The predictive value of low heart rate and heart rate variability during stress for reoffending in delinquent male adolescents. *Psychophysiology*, *48*, 1597–1604. <http://dx.doi.org/10.1111/j.1469-8986.2011.01233.x>
- Diamond, L. M., Fagundes, C. P., & Butterworth, M. R. (2012). Attachment style, vagal tone, and empathy during mother-adolescent interactions. *Journal of Research on Adolescence*, *22*, 165–184. <http://dx.doi.org/10.1111/j.1532-7795.2011.00762.x>
- Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. *Psychological Bulletin*, *130*, 355–391. <http://dx.doi.org/10.1037/0033-2909.130.3.355>
- Egizio, V. B., Jennings, J. R., Christie, I. C., Sheu, L. K., Matthews, K. A., & Gianaros, P. J. (2008). Cardiac vagal activity during psychological stress varies with social functioning in older women. *Psychophysiology*, *45*, 1046–1054. <http://dx.doi.org/10.1111/j.1469-8986.2008.00698.x>
- Eisenberg, N., & Morris, A. S. (2003). Children's emotion-related regulation. In R. V. Kail (Ed.), *Advances in child development and behavior* (Vol. 30, pp. 189–229). San Diego, CA: Academic.
- Eisenberg, N., Sulik, M. J., Spinrad, T. L., Edwards, A., Eggum, N. D., Liew, J., . . . Hart, D. (2012). Differential susceptibility and the early development of aggression: Interactive effects of respiratory sinus arrhythmia and environmental quality. *Developmental Psychology*, *48*, 755–768. <http://dx.doi.org/10.1037/a0026518>
- Ellis, B. J., Essex, M. J., & Boyce, W. T. (2005). Biological sensitivity to context: II. Empirical explorations of an evolutionary-developmental theory. *Development and Psychopathology*, *17*, 303–328. <http://dx.doi.org/10.1017/S0954579405050157>
- El-Sheikh, M. (2005a). Does poor vagal tone exacerbate child maladjustment in the context of parental problem drinking? A longitudinal examination. *Journal of Abnormal Psychology*, *114*, 735–741. <http://dx.doi.org/10.1037/0021-843X.114.4.735>
- El-Sheikh, M. (2005b). Stability of respiratory sinus arrhythmia in children and young adolescents: A longitudinal examination. *Development and Psychopathology*, *46*, 66–74. <http://dx.doi.org/10.1002/dev.20036>
- El-Sheikh, M., Hinnant, J. B. (2011). Marital conflict, respiratory sinus arrhythmia, and allostatic load: Interrelations and associations with the development of children's externalizing behavior. *Development and Psychopathology*, *23*, 815–829. <http://dx.doi.org/10.1017/S0954579411000320>
- El-Sheikh, M., Hinnant, J. B., & Erath, S. (2011). Developmental trajectories of delinquency symptoms in childhood: The role of marital conflict and autonomic nervous system activity. *Journal of Abnormal Psychology*, *120*, 16–32. <http://dx.doi.org/10.1037/a0020626>
- El-Sheikh, M., Kouros, C. D., Erath, S., Cummings, E. M., Keller, P., & Staton, L. (2009). Marital conflict and children's externalizing behavior: Interactions between parasympathetic and sympathetic nervous system activity. *Monographs of the Society for Research in Child Development*, *74*(7), 1–79.
- Fabes, R. A., & Eisenberg, N. (1997). Regulatory control and adults' stress-related responses to daily life events. *Journal of Personality and Social Psychology*, *73*, 1107–1117. <http://dx.doi.org/10.1037/0022-3514.73.5.1107>
- Farrell, A. D., Kung, E. M., White, K. S., & Valois, R. F. (2000). The structure of self-reported aggression, drug use, and delinquent behaviors during early adolescence. *Journal of Clinical Child Psychology*, *29*, 282–292. [http://dx.doi.org/10.1207/S15374424jccp2902\\_13](http://dx.doi.org/10.1207/S15374424jccp2902_13)
- Gentzler, A. L., Santucci, A. K., Kovacs, M., & Fox, N. A. (2009). Respiratory sinus arrhythmia reactivity predicts emotion regulation and depressive symptoms in at-risk and control children. *Biological Psychology*, *82*, 156–163. <http://dx.doi.org/10.1016/j.biopsycho.2009.07.002>
- Goodman, R., & Scott, S. (1999). Comparing the strengths and difficulties questionnaire and the child behavior checklist: Is small beautiful? *Journal of Abnormal Child Psychology*, *27*, 17–24. <http://dx.doi.org/10.1023/A:1022658222914>
- Granic, I., Hollenstein, T., Dishion, T. J., & Patterson, G. R. (2003). Longitudinal analysis of flexibility and reorganization in early adolescence: A dynamic systems study of family interactions. *Developmental Psychology*, *39*, 606–617. <http://dx.doi.org/10.1037/0012-1649.39.3.606>
- Graziano, P., & Derefinko, K. (2013). Cardiac vagal control and children's adaptive functioning: A meta-analysis. *Biological Psychology*, *94*, 22–37. <http://dx.doi.org/10.1016/j.biopsycho.2013.04.011>
- Graziano, P. A., Keane, S. P., & Calkins, S. D. (2007). Cardiac vagal regulation and early peer status. *Child Development*, *78*, 264–278. <http://dx.doi.org/10.1111/j.1467-8624.2007.00996.x>
- Gunnar, M. R., Frenn, K., Wewerka, S. S., & Van Ryzin, M. J. (2009). Moderate versus severe early life stress: Associations with stress reactivity and regulation in 10–12-year-old children. *Psychoneuroendocrinology*, *34*, 62–75. <http://dx.doi.org/10.1016/j.psyneuen.2008.08.013>
- Hastings, P. D., Buss, K. A., & Dennis, T. A. (2012). Introduction to section one: Integrative approaches to the study of physiology and

- emotion. *Monographs of the Society for Research in Child Development*, 77(2), 6–7. <http://dx.doi.org/10.1111/j.1540-5834.2011.00655.x>
- Hastings, P. D., & Miller, J. G. (2014). Autonomic regulation, polyvagal theory, and children's prosocial development. In L. M. Padilla-Walker & G. Carlo (Eds.), *Prosocial development: A multidimensional approach*. New York, NY: Oxford University Press. <http://dx.doi.org/10.1093/acprof:oso/9780199964772.003.0006>
- Hastings, P. D., Nuselovici, J. N., Utendale, W. T., Coutya, J., McShane, K. E., & Sullivan, C. (2008). Applying the polyvagal theory to children's emotion regulation: Social context, socialization, and adjustment. *Biological Psychology*, 79, 299–306. <http://dx.doi.org/10.1016/j.biopsycho.2008.07.005>
- Hinnant, J. B., & El-Sheikh, M. (2009). Children's externalizing and internalizing symptoms over time: The role of individual differences in patterns of RSA responding. *Journal of Abnormal Child Psychology*, 37, 1049–1061. <http://dx.doi.org/10.1007/s10802-009-9341-1>
- Houlberg, B. J., Henry, C. S., & Morris, A. S. (2012). Family interactions, exposure to violence, and emotion regulation: Perceptions of children and early adolescents at risk. *Family Relations*, 61, 283–296. <http://dx.doi.org/10.1111/j.1741-3729.2011.00699.x>
- Ingoldsby, E. M., & Shaw, D. S. (2002). Neighborhood contextual factors and early-starting antisocial pathways. *Clinical Child and Family Psychology Review*, 5, 21–55. <http://dx.doi.org/10.1023/A:1014521724498>
- Katz, L. F., & Rigerink, T. (2012). Domestic violence and emotion socialization. *Monographs of the Society for Research in Child Development*, 77(2), 52–60. <http://dx.doi.org/10.1111/j.1540-5834.2011.00661.x>
- Kok, B. E., & Fredrickson, B. L. (2010). Upward spirals of the heart: Autonomic flexibility, as indexed by vagal tone, reciprocally and prospectively predicts positive emotions and social connectedness. *Biological Psychology*, 85, 432–436. <http://dx.doi.org/10.1016/j.biopsycho.2010.09.005>
- Kop, W. J., Synowski, S. J., Newell, M. E., Schmidt, L. A., Waldstein, S. R., & Fox, N. A. (2011). Autonomic nervous system reactivity to positive and negative mood induction: The role of acute psychological responses and frontal electrocortical activity. *Biological Psychology*, 86, 230–238. <http://dx.doi.org/10.1016/j.biopsycho.2010.12.003>
- Kreibig, S. D. (2010). Autonomic nervous system activity in emotion: A review. *Biological Psychology*, 84, 394–421. <http://dx.doi.org/10.1016/j.biopsycho.2010.03.010>
- Liew, J., Eisenberg, N., Spinrad, T. L., Eggum, N. D., Haugen, R. G., Kupfer, A., . . . Baham, M. E. (2011). Physiological regulation and fearfulness as predictors of young children's empathy-related reactions. *Social Development*, 20, 111–113. <http://dx.doi.org/10.1111/j.1467-9507.2010.00575.x>
- Miller, J. G., Choccol, C., Nuselovici, J. N., Utendale, W. T., Simard, M., & Hastings, P. D. (2013). Children's dynamic RSA change during anger and its relations with parenting, temperament, and control of aggression. *Biological Psychology*, 92, 417–425. <http://dx.doi.org/10.1016/j.biopsycho.2012.12.005>
- MindWare Technologies Ltd. (2010). *MindWare heart rate variability (HRV) analysis user reference guide Version 3.0.9*. [Computer software] Ghanna, OH: MindWare Technologies Ltd.
- Nugent, A. C., Bain, E. E., Thayer, J. F., Sollers, J. J., III, & Drevets, W. C. (2011). Heart rate variability during motor and cognitive tasks in females with major depressive disorder. *Psychiatry Research: Neuroimaging*, 191, 1–8. <http://dx.doi.org/10.1016/j.psychres.2010.08.013>
- Obrovčić, J. (2012). How can the study of physiological reactivity contribute to our understanding of adversity and resilience processes in development? *Development and Psychopathology*, 24, 371–387. <http://dx.doi.org/10.1017/S0954579412000053>
- Obrovčić, J., & Boyce, W. T. (2012). Developmental psychophysiology of emotion processes. *Monographs of the Society for Research in Child Development*, 77(2), 120–128. <http://dx.doi.org/10.1111/j.1540-5834.2011.00670.x>
- Obrovčić, J., Bush, N. R., & Boyce, W. T. (2011). The interactive effect of marital conflict and stress reactivity on externalizing and internalizing symptoms: The role of laboratory stressors. *Development and Psychopathology*, 23, 101–114. <http://dx.doi.org/10.1017/S0954579410000672>
- Obrovčić, J., Bush, N. R., Stamperdahl, J., Adler, N. E., & Boyce, W. T. (2010). Biological sensitivity to context: The interactive effects of stress reactivity and family adversity on socioemotional behavior and school readiness. *Child Development*, 81, 270–289. <http://dx.doi.org/10.1111/j.1467-8624.2009.01394.x>
- Overbeek, T. J., van Boxtel, A., & Westerink, J. H. (2012). Respiratory sinus arrhythmia responses to induced emotional states: Effects of RSA indices, emotion induction method, age, and sex. *Biological Psychology*, 91, 128–141. <http://dx.doi.org/10.1016/j.biopsycho.2012.05.011>
- Porges, S. W. (1995a). Cardiac vagal tone: A physiological index of stress. *Neuroscience and Biobehavioral Reviews*, 19, 225–233. [http://dx.doi.org/10.1016/0149-7634\(94\)00066-A](http://dx.doi.org/10.1016/0149-7634(94)00066-A)
- Porges, S. W. (1995b). Orienting in a defensive world: Mammalian modifications of our evolutionary heritage. A Polyvagal Theory. *Psychophysiology*, 32, 301–318. <http://dx.doi.org/10.1111/j.1469-8986.1995.tb01213.x>
- Porges, S. W. (2003). Social engagement and attachment: A phylogenetic perspective. *Annals of the New York Academy of Sciences*, 1008, 31–47. <http://dx.doi.org/10.1196/annals.1301.004>
- Porges, S. W. (2007). The polyvagal perspective. *Biological Psychology*, 74, 116–143. <http://dx.doi.org/10.1016/j.biopsycho.2006.06.009>
- Porges, S. W. (2011). *The Polyvagal theory: Neurophysiological foundations of emotions, attachment, communication, and self-regulation*. New York, NY: Norton.
- Porges, S. W., Heilman, K. J., Bazhenova, O. V., Bal, E., Doussard-Roosevelt, J. A., & Koledin, M. (2007). Does motor activity during psychophysiological paradigms confound the quantification and interpretation of heart rate and heart rate variability measures in young children? *Developmental Psychobiology*, 49, 485–494. <http://dx.doi.org/10.1002/dev.20228>
- Rash, J. A., & Prkachin, K. M. (2013). Cardiac vagal reactivity during relived sadness is predicted by affect intensity and emotional intelligence. *Biological Psychology*, 92, 106–113. <http://dx.doi.org/10.1016/j.biopsycho.2012.11.009>
- Raudenbush, S. W., Bryk, A. S., & Congdon, R. (2011). *HLM 7 for Windows* [Computer software]. Skokie, IL: Scientific Software International, Inc.
- Rigerink, T., Katz, L. F., & Hessler, D. M. (2010). Domestic violence and longitudinal associations with children's physiological regulation abilities. *Journal of Interpersonal Violence*, 25, 1669–1683. <http://dx.doi.org/10.1177/0886260509354589>
- Ritz, T. (2009). Studying noninvasive indices of vagal control: The need for respiratory control and the problem of target specificity. *Biological Psychology*, 80, 158–168. <http://dx.doi.org/10.1016/j.biopsycho.2008.08.003>
- Steinberg, L. (2001). We know some things: Parent–adolescent relationships in retrospect and prospect. *Journal of Research on Adolescence*, 11, 1–19. <http://dx.doi.org/10.1111/1532-7795.00001>
- Steinberg, L., & Avenevoli, S. (2000). The role of context in the development of psychopathology: A conceptual framework and some speculative propositions. *Child Development*, 71, 66–74. <http://dx.doi.org/10.1111/1467-8624.00119>
- Steinberg, L., & Morris, A. S. (2001). Adolescent development. *Annual Review of Psychology*, 52, 83–110. <http://dx.doi.org/10.1146/annurev.psych.52.1.83>
- Sullivan, T. N., Helms, S. W., Kliever, W., & Goodman, K. L. (2010). Associations between sadness and anger regulation coping, emotional expression, and physical and relational aggression among urban adoles-

- cents. *Social Development*, 19, 30–51. <http://dx.doi.org/10.1111/j.1467-9507.2008.00531.x>
- Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. (1996). Heart rate variability: Standards of measurement, physiological interpretation and clinical use. *Circulation*, 93, 1043–1065. <http://dx.doi.org/10.1161/01.CIR.93.5.1043>
- Thayer, J. F., Hansen, A. L., Saus-Rose, E., & Johnsen, B. H. (2009). Heart rate variability, prefrontal neural function, and cognitive performance: The neurovisceral integration perspective on self-regulation, adaptation, and health. *Annals of Behavioral Medicine*, 37, 141–153. <http://dx.doi.org/10.1007/s12160-009-9101-z>
- Thompson, R. A. (1994). Emotion regulation: A theme in search of definition. *Monographs of the Society for Research in Child Development*, 59(2–3), 25–52. <http://dx.doi.org/10.2307/1166137>
- Tininenko, J. R., Measelle, J. R., Ablow, J. C., & High, R. (2012). Respiratory control when measuring respiratory sinus arrhythmia during a talking task. *Biological Psychology*, 89, 562–569. <http://dx.doi.org/10.1016/j.biopsycho.2011.12.022>
- Vasilev, C. A., Crowell, S. E., Beauchaine, T. P., Mead, H. K., & Gatzke-Kopp, L. M. (2009). Correspondence between physiological and self-report measures of emotion dysregulation: A longitudinal investigation of youth with and without psychopathology. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 50, 1357–1364. <http://dx.doi.org/10.1111/j.1469-7610.2009.02172.x>
- Vögele, C., Sorg, S., Studtmann, M., & Weber, H. (2010). Cardiac autonomic regulation and anger coping in adolescents. *Biological Psychology*, 85, 465–471. <http://dx.doi.org/10.1016/j.biopsycho.2010.09.010>
- Whitson, S. M., & El-Sheikh, M. (2003). Moderators of family conflict and children's adjustment and health. *Journal of Emotional Abuse*, 3, 47–73. [http://dx.doi.org/10.1300/J135v03n01\\_03](http://dx.doi.org/10.1300/J135v03n01_03)
- Zeman, J., Shipman, K., & Penza-Clyve, S. (2001). Development and initial validation of the Children's Sadness Management Scale. *Journal of Nonverbal Behavior*, 25, 187–205. <http://dx.doi.org/10.1023/A:1010623226626>
- Zeman, J., Shipman, K., & Suveg, C. (2002). Anger and sadness regulation: Predictions to internalizing and externalizing symptoms in children. *Journal of Clinical Child and Adolescent Psychology*, 31, 393–398. [http://dx.doi.org/10.1207/S15374424JCCP3103\\_11](http://dx.doi.org/10.1207/S15374424JCCP3103_11)

Received May 20, 2014

Revision received November 2, 2014

Accepted November 4, 2014 ■